SCHEDULING STATUS:

S5

PROPRIETARY NAME (AND DOSAGE FORM): **TRIEMTA TABLETS** (Film-coated Tablets)

COMPOSITION:

Each film-coated tablet contains efavirenz 600 mg, emtricitabine 200 mg, tenofovir disoproxil fumarate 300 mg. Sugar free.

Tablet core: Povidone, microcrystalline cellulose, croscarmellose sodium, hydroxypropyl cellulose, sodium lauryl sulfate, magnesium stearate (vegetable grade), pregelatinised starch, polyvinyl alcohol, titanium dioxide (C.I.No. 77891), polyethylene glycol, talc, red iron oxide (C.I. No. 77491), black iron oxide (C.I. No. 77499).

PHARMACOLOGICAL CLASSIFICATION:

A20.2.8 Antimicrobial (chemotherapeutic) agents. Antiviral agents.

PHARMACOLOGICAL ACTION:

Pharmacodynamic properties:

TRIEMTA TABLETS is a fixed dose combination tablet containing efavirenz, emtricitabine and tenofovir disoproxil fumarate (tenofovir DF). Efavirenz is a non-nucleoside reverse transcriptase inhibitor (NNRTI), emtricitabine is a synthetic nucleoside analogue of cytidine and tenofovir DF is converted in vivo to tenofovir, an acyclic nucleoside phosphonate (nucleotide) analogue of adenosine 5'-monophosphate.

Efavirenz:

Efavirenz is a non-nucleoside reverse transcriptase inhibitor (NNRTI) of HIV-1. Efavirenz activity is mediated predominantly by non-competitive inhibition of HIV-1 reverse transcriptase (RT). Efavirenz does not inhibit HIV-2 RT and human cellular DNA polymerases α, β, γ, and δ.

Emtricitabine:

Emtricitabine, a nucleoside reverse transcriptase inhibitor (NRTI), is a synthetic nucleoside analogue of cytidine. Cellular enzymes phosphorylate emtricitabine to emtricitabine 5'-triphosphate. Emtricitabine 5'-triphosphate competes with the natural substrate, deoxycytidine 5'-triphosphate and is incorporated into nascent viral DNA leading to chain termination. Emtricitabine 5'-triphosphate weakly inhibits mammalian DNA polymerase α, β, ε and mitochondrial DNA polymerase γ.

Tenofovir disoproxil fumarate:

Tenofovir DF, a NRTI, is an acyclic nucleoside phosphonate diester analogue of adenosine monophosphate and requires initial diester hydrolysis for conversion to tenofovir and subsequent phosphorylations by cellular enzymes to form tenofovir diphosphate. Tenofovir diphosphate inhibits the activity of HIV-1 RT through competing with the natural substrate deoxyadenosine 5'-triphosphate. After incorporation into DNA, the DNA chain is terminated. Tenofovir diphosphate weakly inhibits mammalian DNA polymerases α, β, and mitochondrial DNA polymerase γ.

Antiviral activity:

Efavirenz, emtricitabine and tenofovir disoproxil fumarate:

In combination studies that evaluated the antiviral activity *in vitro* of emtricitabine and efavirenz against laboratory and tenofovir together and emtricitabine and tenofovir together, additive to synergistic antiviral effects were observed.

Resistance:

Efavirenz, emtricitabine and tenofovir disoproxil fumarate:

HIV-1 isolates demonstrating reduced susceptibility to the emtricitabine and tenofovir combination have been selected *in vitro*.

Following genotypic analysis, the M184V/I and/or K65R amino acid substitutions were identified in the viral RT.

Efavirenz:

Clinical isolates with reduced susceptibility *in vitro* to efavirenz have been obtained. The most frequently observed amino acid substitution in clinical studies with efavirenz is K103N (54 %). One or more RT substitutions at amino acid positions 98, 100, 101, 103, 106, 108, 188, 190, 225, 227, and 230 were observed in patients failing treatment with efavirenz in combination with other antiretrovirals. Other resistance substitutions observed were commonly included L100I (7 %), K101E/Q/R (14 %), V108I (11 %), G190S/T/A (7 %), P225H (18 %), and M230I/L (11 %).

HIV-1 isolates with reduced susceptibility to efavirenz (more than 350-fold increase in EC90 value) emerged rapidly under selection *in vitro*. Genotypic characterisation of these viruses identified mutations resulting in single amino acid substitutions L100I or V179D, double substitutions L100I/V108I, and triple substitutions L100I/V179D/Y181C in RT.

Emtricitabine:

HIV isolates resistant to emtricitabine have been selected *in vitro*. Genotypic analysis of these viruses showed point mutations that reduced emtricitabine susceptibility was associated with a mutation in the HIV RT codon M184. This mutation caused an amino acid substitution of methionine by valine or isoleucine (M184V/I).

Tenofovir disoproxil fumarate:

Isolates of HIV-1 that exhibited reduced susceptibility to tenofovir have been selected *in vitro*. These viruses expressed a K65R mutation in reverse transcriptase and demonstrated a reduced susceptibility to tenofovir of between 2- and 4-fold.

Cross-resistance:

Efavirenz, emtricitabine and tenofovir disoproxil fumarate:

Cross-resistance has been recognised among NNRTIs. It is also known that cross-resistance may develop among certain NRTIs. The M184V/I and/or K65R substitutions selected *in vitro* by the combination of emtricitabine and tenofovir are also seen in some HIV isolates from individuals failing treatment with tenofovir in combination with either lamivudine or emtricitabine, and either abacavir or didanosine. Combinations of these medicines may therefore develop in patients whose virus harbours either or both of these amino acid substitutions.

Pharmacokinetic properties:

Efavirenz:

Absorption:

Peak efavirenz plasma concentrations of 1.6 – 9.1 μM are reached by 5 hours following single oral doses of 100 mg to 1600 mg given to uninfected volunteers. Although dose-related increases in C_{max} and AUC are seen for doses up to 1600 mg, these increases are less than proportional, indicating diminished absorption at higher doses. Steady state plasma concentrations are reached in 6 to 10 days.

Distribution:

Efavirenz is highly plasma protein bound (approximately 99,5 – 99,75 %) and predominantly to albumin. Cerebrospinal fluid concentrations ranged from 0,26 to 1,19 % (mean 0,69 %) of the corresponding plasma concentrations in HIV-1 infected individuals administered 200 to 600 mg efavirenz once daily for at least one month. This is about three times higher than the non-protein-bound (free) fraction of efavirenz in plasma.

Metabolism:

Efavirenz is principally metabolised by the cytochrome P450 system to hydroxylated metabolites with subsequent glucuronidation of these hydroxylated metabolites. The metabolites are not active against HIV-1. CYP3A4 and CYP2B6 are the major isozymes involved in efavirenz metabolism.

Efavirenz, through induction of P450 enzymes, induces its own metabolism.

Elimination:

Efavirenz has a long terminal half-life of 52 – 76 hours and 40 – 55 hours after single and multiple doses, respectively. Subsequent to the administration of a radio-labelled dose, 14 – 34 % of efavirenz is recovered in the urine (mostly as metabolites) and 16 – 61 % is recovered in the faeces (mostly as parent medicine).

Emtricitabine:

The pharmacokinetic properties of emtricitabine are summarised in Table 1.

Absorption: Following oral administration of emtricitabine (200 mg), emtricitabine is rapidly absorbed with peak plasma concentrations occurring at 1 – 2 hours post-dose.

Distribution: *In vitro* binding of emtricitabine to human plasma proteins is < 4 % and is independent of concentration over the range of 0,02 – 200 μg/ml.

Metabolism and elimination: Following administration of radio-labelled emtricitabine, approximately 86 % is recovered in the urine and 13 % is recovered as metabolites. The metabolites of emtricitabine include 2'-sulfoxide diastereomers and their glucuronic acid conjugates. Emtricitabine is eliminated by a combination of glomerular filtration and active tubular secretion. Following a single oral dose of emtricitabine (200 mg), the plasma emtricitabine half-life is approximately 10 hours.

Tenofovir Disoproxil Fumarate:

The pharmacokinetic properties of tenofovir disoproxil fumarate are summarised in Table 1.

Absorption: Following oral administration of tenofovir, maximum tenofovir serum concentrations are achieved in 1,0 ± 0,4 hour.

Distribution: *In vitro* binding of tenofovir to human plasma proteins < 0,7 % is independent of concentration over the range of 0,01 – 25 μg/ml.

Metabolism and elimination: Approximately 70 – 80 % of the intravenous dose of tenofovir is recovered as unchanged in the urine. Tenofovir is eliminated by a combination of glomerular filtration and active tubular secretion. Following a single oral dose of tenofovir, the terminal elimination half-life of tenofovir is approximately 17 hours.

Table 1: Single Dose Pharmacokinetic Parameters for Emtricitabine and Tenofovir in Adults

	Emtricitabine	Tenofovir
Fasted Oral Bioavailability[1] (%)	92 (83,1 – 106,4)	25 (NC[2] – 45,0)
Plasma Terminal Elimination Half-Life[3] (hr)	10 (7,4 – 18,0)	17 (12,0 – 25,7)
C_{max}[3] (μg/ml)	1,8 ± 0,72[4]	0,30 ± 0,09
AUC[3] (μg•hr/ml)	10,0 ± 3,12[4]	2,29 ± 0,69
CL/F[3] (ml/min)	302 ± 94	1043 ± 115
CL_{renal}[3] (ml/min)	213 ± 89	243 ± 33

[1] NC = Not calculated

[2] Median (range)

[3] Mean (± SD)

[4] Data presented as steady state values.

Effects of food on oral absorption:

TRIEMTA TABLETS has not been evaluated in the presence of food. Administration of efavirenz tablets with a high fat meal increased the mean AUC and C_{max} of efavirenz by 28 % and 79 % respectively, compared to administration in the fasted state. Compared to fasted administration, dosing of tenofovir DF and emtricitabine in combination with either a high fat meal or a light meal increased the mean AUC and C_{max} of tenofovir by 35 % and 15 % respectively, without affecting emtricitabine exposures.

Special populations:

Paediatric and elderly patients

Pharmacokinetic studies of tenofovir DF have not been performed in paediatric patients (less than 18 years). Efavirenz has not been studied in paediatric patients below 3 years of age or who weigh less than 13 kg. Emtricitabine has been studied in paediatric patients from 3 months to 17 years of age. **TRIEMTA TABLETS** is not recommended for paediatric administration. Pharmacokinetics of efavirenz, emtricitabine, and tenofovir have not been fully evaluated in the elderly (more than 65 years) (see **WARNINGS AND SPECIAL PRECAUTIONS:**).

Patients with impaired renal function

Efavirenz: The pharmacokinetics of efavirenz have not been studied in patients with renal insufficiency, however less than 1 % of efavirenz is excreted unchanged in the urine, so the impact of renal impairment on efavirenz elimination should be minimal.

Emtricitabine and tenofovir disoproxil fumarate: The pharmacokinetics of emtricitabine and tenofovir DF are altered in patients with renal impairment. In patients with creatinine clearance less than 50 ml/min, C_{max} and AUC_{∞} of emtricitabine and tenofovir were increased (see **CONTRAINDICATIONS, WARNINGS AND SPECIAL PRECAUTIONS, Renal impairment**).

Patients with hepatic impairment

Efavirenz: The pharmacokinetics of efavirenz have not been adequately studied in patients with hepatic impairment (see **WARNINGS AND SPECIAL PRECAUTIONS:**).

Emtricitabine: The pharmacokinetics of emtricitabine have not been studied in patients with hepatic impairment. However, emtricitabine is not significantly metabolised by liver enzymes, so the impact of liver impairment should be limited.

Tenofovir disoproxil fumarate: The pharmacokinetics of tenofovir following a 300 mg dose of tenofovir DF have been studied in non-HIV infected patients with moderate to severe hepatic impairment. There were no substantial alterations in tenofovir pharmacokinetics in patients with hepatic impairment compared with unimpaired patients.

INDICATIONS:

TRIEMTA TABLETS is indicated as mono- or combination therapy with other anti-retroviral medicines for HIV-1 infection in adults.

CONTRAINDICATIONS:

TRIEMTA TABLETS is contraindicated in patients:

- With previously demonstrated hypersensitivity to efavirenz, emtricitabine or tenofovir or to any of the other components of **TRIEMTA TABLETS**.
- Severe hepatic impairment (Child Pugh Class C).
- A history of previous liver injury/failure with efavirenz containing antiretroviral treatment (ART).
- With moderate to severe impairment of renal function (creatine clearance < 50 ml) (see **WARNINGS AND SPECIAL PRECAUTIONS** and **DOSAGE AND DIRECTIONS FOR USE**).
- Who are concurrently receiving treatment with cisapride, midazolam, triazolam, pimozide, bepridil or ergot alkaloids (for example ergotamine, dihydroergotamine, ergonovine and methylergonovine), because competition for CYP3A4 by the efavirenz in **TRIEMTA TABLETS** could result in inhibition of the metabolism of these medicines and create the potential for serious and/or life-threatening adverse events (e.g., cardiac dysrhythmias, prolonged sedation or respiratory depression).
- Who are concurrently using voriconazole, since efavirenz significantly decreases voriconazole plasma concentrations, while voriconazole significantly increases efavirenz plasma concentrations. (See **INTERACTIONS**). Since **TRIEMTA TABLETS** is a fixed-dose combination tablet, the dosage of voriconazole cannot be altered; therefore voriconazole and **TRIEMTA TABLETS** cannot be co-administered.
- Who are pregnant or breastfeeding (see **PREGNANCY AND LACTATION**).
- Who are concurrently using St. John's Wort (*Hypericum perforatum*) or St. John's Wort containing medicines (see **INTERACTIONS**).
- **TRIEMTA TABLETS** is not recommended for use in children and adolescents below 18 years of age due to lack of data on safety and efficacy.

WARNINGS AND SPECIAL PRECAUTIONS:

LACTIC ACIDOSIS AND SEVERE HEPATOMEGALY WITH STEATOSIS, INCLUDING FATAL CASES, HAVE BEEN REPORTED WITH THE USE OF NUCLEOSIDE ANALOGUES ALONE OR IN COMBINATION WITH OTHER ANTI-RETROVIRALS.

TRIEMTA TABLETS ARE NOT INDICATED FOR THE TREATMENT OF CHRONIC HBV INFECTION AND SAFETY AND EFFICACY OF **TRIEMTA TABLETS** HAVE NOT BEEN ESTABLISHED IN PATIENTS CO-INFECTED WITH HBV AND HIV. SEVERE ACUTE EXACERBATIONS OF HEPATITIS B HAVE BEEN REPORTED IN PATIENTS WHO HAVE BEEN CO-INFECTED WITH HBV AND HIV AND HAVE STOPPED TREATMENT WITH **TRIEMTA TABLETS**. WHETHER OR NOT EMTRICITABINE, THE EXACERBATION OF HEPATITIS B MAY BE ASSOCIATED WITH LIVER DECOMPENSATION AND LIVER FAILURE. HEPATIC FUNCTION SHOULD BE MONITORED CLOSELY WITH BOTH CLINICAL AND LABORATORY FOLLOW-UP FOR AT LEAST SEVERAL MONTHS IN PATIENTS CO-INFECTED WITH HIV AND HBV WHO HAVE STOPPED **TRIEMTA TABLETS**. IF APPROPRIATE, INITIATION OF ANTI-HEPATITIS B THERAPY MAY BE WARRANTED.

Lactic acidosis / hyperlactataemia:

Use of **TRIEMTA TABLETS** can result in potentially fatal lactic acidosis as a consequence of mitochondrial dysfunction.

Clinical findings are non-specific, and include nausea, vomiting, abdominal pain, dyspnoea, fatigue and weakness.

In patients with precipitous symptoms, measure the venous lactate level (normal < 2 mmol/l) and the serum bicarbonate and respiratory rate as follows:

- **Lactate 2 – 5 mmol/l with minimum symptoms:** switch to medicines that are less likely to cause lactic acidosis.
- **Lactate 5 – 10 mmol/l with symptoms and or with reduced bicarbonate:** Stop NRTIs and change treatment urgently. Once lactate has been stabilised, re-introduce these NRTIs if lactic acidosis. Exclude other causes, (e.g. sepsis, uraemia, diabetic ketoacidosis, thyrotoxicosis and hyperthyroidism).
- **Lactate > 10 mmol/l: STOP all therapy** (80 % mortality).

The above lactate values may not be applicable to children patients.

Caution should be exercised when administering **TRIEMTA TABLETS** to patients with known factors for liver disease.

Treatment with **TRIEMTA TABLETS** should be suspended in any patient who develops clinical or laboratory findings suggestive of lactic acidosis or hepatotoxicity.

Mitochondrial dysfunction:

Nucleoside and nucleotide analogues have been demonstrated *in vitro* and *in vivo* to cause a variable degree of mitochondrial damage. There have been reports of mitochondrial dysfunction in HIV-negative infants exposed *in utero* and/or post-natally to nucleoside analogues. Apart from lactic acidosis/hyperlactataemia (see above) other manifestations of mitochondrial dysfunction include haematological disorder (anaemia, neutropenia), pancytopenia, myopathy, lipodystrophy, and onset neurological disorders have been reported (hypertonia, convulsion, abnormal behaviour). It is not known whether the neurological disorders are transient or permanent. Any child exposed *in utero* to nucleoside and nucleotide analogues, even HIV negative infants/children, should have clinical and laboratory follow-up and should be fully investigated for possible mitochondrial dysfunction in case of relevant signs or symptoms.

Pancreatitis:

Pancreatitis has been observed in some patients receiving **TRIEMTA TABLETS**. Pancreatitis must be considered whenever a patient develops clinical symptoms, or laboratory or elevated biochemical markers. Discontinue use of **TRIEMTA TABLETS** until diagnosis of pancreatitis is excluded.

Liver disease:

Use of **TRIEMTA TABLETS** can result in hepatotoxicity due to non-alcoholic fatty liver disease (hepatic steatosis). The safety and efficacy of **TRIEMTA TABLETS** have been established in patients with significant underlying liver disorders/diseases. In case of concomitant antiviral therapy for hepatitis B or C, please also refer to the relevant package inserts for these medicines.

Patients with pre-existing liver dysfunction including chronic active hepatitis have an increased frequency of liver function abnormalities during combination antiretroviral therapy and should be monitored. If there is any evidence of worsening liver disease in such patients, temporary or permanent discontinuation of treatment must be considered.

There is some evidence that efavirenz is associated with three clinical pathological patterns of drug induced liver failure in HIV positive patients of which the sub massive necrosis histological pattern seems to be associated with a high morbidity/mortality risk and may present many months after therapy has been initiated or even stopped. Risk factors include younger age, CD4+ counts > 350 cells/μl and female gender.

Early detection and treatment of the liver failure and the immediate discontinuation of **TRIEMTA TABLETS** or efavirenz containing medicines should be stressed. Patients who discontinued treatment with **TRIEMTA TABLETS** should be followed up for symptoms/signs of liver failure for up to 12 months.

TRIEMTA TABLETS is not recommended in patients with moderate to severe hepatic impairment because there could be insufficient data to determine whether dose adjustments are required.

Liver enzymes:

Monitoring of liver enzyme levels is recommended in patients with known or suspected history of hepatitis B or C infection and in patients treated with other medicines associated with hepatic toxicity. The benefit-risk ratio of continued therapy with **TRIEMTA TABLETS** has to be weighed in the unknown risks of significant hepatic toxicity in patients who show persistent elevations of serum transaminases to greater than 5 times the upper limit of the normal range (see **Side Effects**).

Patients on **TRIEMTA TABLETS** or efavirenz containing antiretroviral treatment (ART) should be regularly monitored for jaundice (including a laboratory bilirubin and liver enzymes) and bleeding tendencies.

Patients with HIV and hepatitis B or C virus co-infection:

It is recommended that all patients co-infected with HIV be tested for the presence of chronic HBV before initiating anti-retroviral therapy. **TRIEMTA TABLETS** are not indicated for the treatment of chronic HBV infection and safety and efficacy of **TRIEMTA TABLETS** have not been established in patients co-infected with HBV and HIV. Severe acute exacerbations of hepatitis B have been reported in patients co-infected with HBV and HIV and who have discontinued **TRIEMTA TABLETS**. In some of these patients treated with emtricitabine, the exacerbation of hepatitis B were associated with liver decompensation and liver failure. Hepatic function should be monitored closely with both clinical and laboratory follow-up for at least several months in patients co-infected with HBV and HIV and who have discontinued **TRIEMTA TABLETS**. If appropriate, initiation of anti-hepatitis B therapy may be warranted.

Patients with chronic hepatitis B or C and treated with antiretroviral therapy are at an increased risk for severe and potentially fatal hepatic adverse reactions.

Medical practitioners should refer to current HIV treatment guidelines for the optimal management of HIV infection in patients co-infected with hepatitis B virus (HBV).

In case of concomitant antiviral therapy for hepatitis B or C, please refer also to the relevant package inserts for these medicines.

In patients with advanced liver disease or cirrhosis, treatment discontinuation is not recommended since post-treatment exacerbation of hepatitis may lead to hepatic decompensation.

Discontinuation of **TRIEMTA TABLETS** therapy in patients co-infected with HIV and HBV may be associated with severe acute exacerbations of hepatitis.

Co-administration with related medicines:

Do not co-administer any other medicine in combination with **TRIEMTA TABLETS** include emtricitabine, tenofovir DF, emtricitabine/tenofovir DF and efavirenz which contain the same active components as **TRIEMTA TABLETS**. Due to similarities between emtricitabine and lamivudine, **TRIEMTA TABLETS** should not be co-administered with medicines containing lamivudine, including lamivudine/zidovudine, lamivudine, and abacavir.

St. John's Wort:

Concomitant use of **TRIEMTA TABLETS** and St. John's Wort (*Hypericum perforatum*) or St. John's Wort containing products is contraindicated. Co-administration of NNRTIs, including efavirenz with St. John's Wort is expected to substantially decrease efavirenz concentrations and may result in suboptimal levels of efavirenz and lead to loss of virologic response and possible resistance to efavirenz or to the class of NNRTIs.

Psychiatric symptoms:

Serious psychiatric adverse events have been reported in patients treated with efavirenz. These side effects include: severe depression, suicidal ideation, nonfatal suicide attempts, aggressive behaviour, paranoid reactions and manic reactions. Treatment with efavirenz is associated with an increase in the occurrence of these psychiatric symptoms.

There have also been rare post-marketing reports of death by suicide, delusions, and psychosis-like behaviour; a causal relationship to the use of efavirenz cannot be determined from these reports.

Patients with serious psychiatric adverse experiences should seek immediate medical evaluation to assess the possibility that the symptoms may be related to the use of efavirenz, therefore discontinuation should be considered.

Nervous system and psychiatric symptoms:

These symptoms included dizziness, insomnia, impaired concentration, somnolence, abnormal dreams and hallucinations. Other reported symptoms are euphoria, agitation, confusion, agitation, amnesia, abnormal thinking, and depersonalization. These symptoms usually begin during the first or second day of therapy and may resolve after the first 2 - 4 weeks of therapy. Patients should be informed that these common symptoms may improve with continued therapy and may not be predictive of subsequent onset of the less frequent psychiatric symptoms. Dosing at bedtime may improve the tolerability of these nervous system symptoms.

Patients receiving **TRIEMTA TABLETS** should be alerted to the potential for additive central nervous system effects when **TRIEMTA TABLETS** is used concomitantly with alcohol or psychoactive medicines.

Renal impairment:

Emtricitabine and tenofovir, two of the active ingredients in **TRIEMTA TABLETS**, are principally eliminated by the kidney. **TRIEMTA TABLETS** should not be administered to patients with creatinine clearance < 50 ml/min (see **Contraindications**).

There have been reports of renal impairment including cases of acute renal failure and Fanconi syndrome (renal tubular injury with severe hypophosphataemia) in association with the use of tenofovir DF (see **Side Effects**).

It is recommended that creatinine clearance be calculated in all patients prior to initiating therapy and as clinically appropriate during therapy with **TRIEMTA TABLETS** (see **Dosage and Directions for Use**). Routine monitoring of calculated creatinine clearance and serum phosphorous should be performed in patients at risk for renal impairment.

The use of **TRIEMTA TABLETS** should be avoided in patients who are concurrently taking or who have recently taken a nephrotoxic medicine.

Skin rash:

Mild-to-moderate rash, which may resolve with continued treatment, has been observed with efavirenz use. Treatment with appropriate antihistamines and/or corticosteroids may improve tolerability and hasten resolution of symptoms. Treatment with **TRIEMTA TABLETS** should be discontinued in patients who experience severe rash associated with blistering, desquamation, mucosal involvement or fever (see **Side effects**).

Cholesterol:

Monitoring of cholesterol and triglyceride levels should be considered in patients treated with **TRIEMTA TABLETS** (see **Side Effects**).

Bone effects:

TRIEMTA TABLETS decreases bone mineral density. Bone monitoring should be considered for HIV infected patients who have a history of pathologic bone fracture or are at risk for osteopenia. Although the effect of supplementation with calcium and vitamin D was not studied, such supplementation may be beneficial for all patients. If bone abnormalities are suspected then appropriate consultation should be obtained.

Osteonecrosis:

Although the aetiology is considered to be multifactorial (including corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index), cases of osteonecrosis have been reported, particularly in patients with advanced HIV-disease and/or long-term exposure to combination antiretroviral therapy (cART). Patients should be advised to seek medical advice if they experience joint aches and pains, joint stiffness or difficulty in movement.

Convulsions:

Convulsions have been observed in patients receiving efavirenz, generally in the presence of known medical history of seizures. Caution must be taken in any patient with a history of seizures.

Patients who are concurrently receiving anticonvulsant medicines are mainly metabolised by the liver, such as phenytoin and phenobarbitone, may require periodic monitoring of plasma levels (see **INTERACTIONS**).

Lipodystrophy and metabolic abnormalities:

Combination antiretroviral therapy has been associated with the redistribution/accumulation of body fat, including central obesity, dorso-cervical fat enlargement (buffalo hump), peripheral wasting, facial wasting, breast enlargement, and elevated serum lipid and glucose levels in HIV Positive patients. Clinical examinations should include evaluation for physical signs of fat redistribution. Patients with evidence of lipodystrophy should have a thorough cardiovascular risk assessment.

Immune Reconstitution Inflammatory Syndrome:

Immune reconstitution inflammatory syndrome (IRIS) is an immunopathological response resulting from rapid restoration of pathogen-specific immune responses to pre-existing antigens combined with immune dysregulation, which occurs shortly after starting combination Anti-Retroviral Therapy (cART). Typically it presents by paradoxical deterioration of opportunistic infections being treated or after with unmasking of an asymptomatic opportunistic disease, often with an atypical inflammatory presentation. IRIS usually develops within the first three months of initiation of cART. Most commonly reported examples are Kaposi's sarcoma, CMV retinitis, mycobacterial infections or reactions to opportunistic diseases are tuberculosis, PCP, cryptococcal meningitis and cryptococcal meningitis. Appropriate treatment of the opportunistic disease should be instituted or continued and cART continued. Inflammatory manifestations generally subside after a few weeks. Severe cases may respond to glucocorticoids, but there is only limited evidence for this in patients with tuberculosis IRIS. Autoimmune disorders (such as Graves' disease) have also been reported as IRIS reactions; however, the reported time to onset is more variable and these events can occur many months after initiation of treatment.

Opportunistic infections:

Patients receiving **TRIEMTA TABLETS** should be advised that they may continue to develop opportunistic infections and other complications of HIV infection, and therefore they should remain under close observation by healthcare professionals experienced in the treatment of patients with associated HIV disease. Regular monitoring of CD4 counts is recommended.

The risk of HIV transmission to others:

Patients should be advised that current antiretroviral therapy, including **TRIEMTA TABLETS**, does not prevent the risk of transmission of HIV to others through sexual contact or blood contamination. Appropriate precautions should continue to be employed.

Paediatric and adolescent use:

The safety and efficacy of **TRIEMTA TABLETS** in children and adolescents less than 18 years of age have not been established, therefore **TRIEMTA TABLETS** is not recommended in this age group.

Use in the elderly:

Clinical studies of fixed dose combinations of emtricitabine and tenofovir, as in **TRIEMTA TABLETS**, did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Generally, caution is required when doses are selected for elderly patients, because of the greater frequency of decreased liver, kidney and cardiac function, and of concomitant disease or therapy with other medicines.

Effects on the ability to drive and use machines:

TRIEMTA TABLETS may lead to dizziness, impaired concentration, and/or drowsiness. Patients should be instructed that, if they develop these symptoms, they should keep away from potentially hazardous tasks, such as driving or operating machinery.

INTERACTIONS:

Efavirenz: Efavirenz has been shown *in vivo* to induce CYP3A4 and CYP2B6. Other compounds that are substrates of CYP3A4 may have decreased plasma concentrations when co-administered with efavirenz. *In vitro* studies have demonstrated that efavirenz inhibits CYP3A4 isozymes in the range of observed efavirenz plasma concentrations. Co-administration of efavirenz with medicines primarily metabolised by these isozymes may result in plasma concentrations that exceed the co-administered medicine. Therefore, appropriate dose adjustments may be necessary for these medicines.

Medicines that induce CYP3A4 activity (e.g., phenobarbitone, rifampicin, rifabutin) would be expected to increase the clearance of efavirenz, resulting in decreased efavirenz concentrations. Efavirenz exposure may be increased by medicines (such as ritonavir) or grapefruit juice which inhibit CYP3A4 (see **Side Effects**).

Emtricitabine and tenofovir disoproxil fumarate: Since emtricitabine and tenofovir are primarily eliminated by the kidneys, co-administration of **TRIEMTA TABLETS** with medicines that reduce renal function or compete for active tubular secretion may increase serum concentrations of emtricitabine, tenofovir, and/or other renally eliminated medicines. Some examples include, but are not limited to: acyclovir, adefovir dipivoxil, cidofovir, ganciclovir, valacyclovir, valganciclovir, aminoglycosides, amphotericin B, foscarnet, pentamidine and interleukin-2.

Co-administration of tenofovir DF and didanosine should be undertaken with caution and patients receiving this combination should be monitored closely for didanosine-associated adverse events. Didanosine should be discontinued in patients who develop didanosine-associated adverse events (for didanosine dosing adjustment recommendations, see Table 3). Suppression of CD4 cell counts has been observed in patients receiving tenofovir DF with didanosine 400 mg daily.

Atazanavir and lopinavir/ritonavir have been shown to increase tenofovir concentrations. The mechanism of this interaction is unknown. Higher tenofovir concentrations could potentiate tenofovir-associated adverse events, including renal disorders. Patients receiving either atazanavir or lopinavir/ritonavir with tenofovir DF should be monitored for tenofovir-associated adverse events. **TRIEMTA TABLETS** should be discontinued in patients who develop tenofovir-associated adverse events (For atazanavir dosing adjustment recommendations see Table 3).

Other important medicine interactions for **TRIEMTA TABLETS** are summarised in Table 2 and 3. The medicine interactions described are based on studies conducted with efavirenz, emtricitabine, tenofovir DF, or emtricitabine/tenofovir DF, or are potential medicine interactions. No medicine interaction studies have been conducted using the combination of actives such as in **TRIEMTA TABLETS**. The tables include potentially significant interactions, but are not all inclusive.

Table: 2

Medicines that are contraindicated or not recommended for use with TRIEMTA TABLETS

Medicine Class: Medicine Name	Clinical Comment
Antifungal: voriconazole	CONTRAINDICATED because efavirenz significantly decreases voriconazole plasma concentrations and co-administration may decrease the therapeutic effectiveness of voriconazole. Also, voriconazole significantly increases efavirenz plasma concentrations, which may increase the risk of efavirenz-associated side effects.
Anti-migraine: ergot derivatives (dihydroergotamine, ergonovine, ergotamine, methylergonovine)	CONTRAINDICATED due to potential for serious and/or life-threatening reactions such as acute ergot toxicity characterised by peripheral vasospasm and ischaemia of the extremities and central nervous system.
Anti-retroviral: efavirenz, emtricitabine, tenofovir DF, lamivudine	Not for use with **TRIEMTA TABLETS** because the active ingredients- emtricitabine, tenofovir DF, emtricitabine/ tenofovir DF and efavirenz are components of **TRIEMTA TABLETS**. Lamivudine is similar to emtricitabine.
Benzodiazepines: midazolam, triazolam	CONTRAINDICATED due to serious and/or life-threatening reactions such as prolonged or increased sedation or respiratory depression.
Calcium channel blocker: bepridil	CONTRAINDICATED due to potential for serious and/or life-threatening reactions such as cardiac dysrhythmias.
GI motility medicine: cisapride	CONTRAINDICATED due to potential for serious and/or life-threatening reactions such as cardiac dysrhythmias.
Neuroleptic: pimozide	CONTRAINDICATED due to potential for serious and/or life-threatening reactions such as cardiac dysrhythmias.
St. John's Wort (*Hypericum perforatum*)	CONTRAINDICATED: Expected to substantially decrease efavirenz, tenofovir, has not been studied in combination with efavirenz.

Established and Other Potentially Significant Medicine Interactions: Alteration in Dose or Regimen May Be Recommended Based on Medicine Interaction Trials or Predicted Interaction

Concomitant Medicine Class: Medicine Name	Effect	Clinical Comment
Antiretroviral medicines		
Protease inhibitor: Amprenavir	↓ amprenavir concentration	Efavirenz has the potential to decrease serum concentrations of amprenavir.
Protease inhibitor: Fosamprenavir calcium	↓ amprenavir concentration	Fosamprenavir is co-administered: Appropriate doses of fosamprenavir and ritonavir have not been established with respect to safety and efficacy have not been established. Fosamprenavir/ritonavir: An additional 100 mg/day (300 mg total) of ritonavir is recommended when **TRIEMTA TABLETS** are administered with fosamprenavir once daily. No change in the ritonavir dose is required when **TRIEMTA TABLETS** is administered with fosamprenavir/ritonavir once daily. No change in the ritonavir dose is required when **TRIEMTA TABLETS** is administered with fosamprenavir/ritonavir twice daily.
Protease inhibitor: Atazanavir	↓ atazanavir concentration ↑ tenofovir concentration	Plasma concentration of atazanavir was decreased by both efavirenz or tenofovir DF. These data are not available to make a dosage recommendation for atazanavir or atazanavir/ritonavir with **TRIEMTA TABLETS**. Therefore co-administration of **TRIEMTA TABLETS** and atazanavir is not recommended due to concerns regarding decreased atazanavir concentrations.
Protease inhibitor: Indinavir	↓ indinavir concentration.	The optimal dose of indinavir, when given in combination with efavirenz, is not known. Increasing the indinavir dose to 1000 mg every 8 hours does not compensate for the increased indinavir metabolism due to efavirenz.
Protease inhibitor: Lopinavir/ritonavir	↓ lopinavir concentration ↑ tenofovir concentration	A dose increase of lopinavir/ritonavir to 600/150 mg (3 tablets) twice daily may be considered when used in combination with efavirenz in treatment-experienced patients where decreased susceptibility to lopinavir is clinically suspected (by treatment history or resistance testing). **Medicines should be monitored for tenofovir-associated adverse events. TRIEMTA TABLETS should be discontinued in patients who develop tenofovir-associated adverse events.**
Protease inhibitor: Ritonavir	↑ ritonavir concentration ↑ efavirenz concentration	When ritonavir 500 mg every 12 hours was co-administered with efavirenz 600 mg once daily, the combination was associated with a higher frequency of adverse clinical experiences (e.g., dizziness, nausea, paraesthesia, elevated liver enzymes). Monitoring of liver enzymes is recommended when **TRIEMTA TABLETS** is used in combination with ritonavir.
Protease Inhibitor: Saquinavir	↑ saquinavir concentration	Should not be used as sole protease inhibitor in combination with **TRIEMTA TABLETS**.
NRTI: Didanosine	↑ didanosine concentration	Higher didanosine concentrations could potentiate didanosine-associated adverse events, including pancreatitis and lactic acidosis. Clinical examinations should include evaluation for physical signs of fat redistribution. Patients with evidence of lipodystrophy should have a thorough cardiovascular risk assessment.
Other medicines		
Anticoagulant: Warfarin	↑ or ↓ warfarin concentration	Plasma concentrations and effects potentially increased or decreased by efavirenz.
Anticonvulsants: Carbamazepine	↓ carbamazepine concentration ↓ efavirenz concentration	There are insufficient data to make a dose recommendation for **TRIEMTA TABLETS**. Alternative anticonvulsant treatment should be used.
Phenytoin Phenobarbitone	↓ anticonvulsant concentration ↓ efavirenz concentration	Potential for reduction in anticonvulsant and/or efavirenz exposure.
Valproic acid/Efavirenz	No clinically significant effect on efavirenz pharmacokinetics. Limited data suggest that there is no clinically significant effect on valproic acid pharmacokinetics.	**TRIEMTA TABLETS** and valproic acid can be co-administered without dose adjustment. Patients should be monitored for seizure control.
Vigabatrin/Efavirenz Gabapentin/Efavirenz	Interaction not studied. Clinically significant interactions are not expected since vigabatrin and gabapentin are exclusively eliminated unchanged in the urine (95 % and 100 %, respectively), compared to the same metabolic enzymes and elimination pathways as efavirenz.	**TRIEMTA TABLETS** and vigabatrin or gabapentin can be co-administered without dose adjustment.
Antidepressant : Sertraline,	↓ sertraline concentration	Increases in sertraline dose should be guided by clinical response.
Paroxetine/Efavirenz	Paroxetine: AUC: ↔ C_{max}: ↔ C_{min}: ↔ Efavirenz: AUC: ↔ C_{max}: ↔ C_{min}: ↔	**TRIEMTA TABLETS** and paroxetine can be co-administered without dosage adjustment.
Fluoxetine/Efavirenz	Interaction not studied. Since fluoxetine shares a similar metab

SKEDUELERINGSTATUS:

S4

EENDOMSNAAM (en ooseervorm):
TRIEMTA TABLETS (Filmbedekte tablette).

SAMESTELLING:
Elke filmbedekte tablet bevat efavirens 600 mg, emtrisitabien 200 mg en tenofovir disoproksielfumaraat 300 mg. Sukkervry.
Ander bestanddele sluit mikrokristallyne sellulose, kroskarmellose natrium, hidroksipropielsellulose, natriumbouteriusftaat, magnesiumsteraat (plantaardig), pigmentgeïnkorporeerde gelatienlaag (hipromelloos), titaandioksied (C.I. 77891), poliëtileenglikol, talk, rooi ysteroksied (C.I. 77491) en swart ysteroksied (C.I. 77499) in.

FARMAKOLOGIESE KLASSIFIKASIE:
A.20.2.8 Antimikrobiese (chemoterapeutiese) middels. Antivirusmiddels.

FARMAKOLOGIESE WERKING:
Farmakodinamiese eienskappe:
TRIEMTA TABLETS is 'n vastedosis kombinasietablet wat efavirens, emtrisitabien en tenofovir disoproksielfumaraat (tenofovir DF) bevat. Efavirens is 'n nie-nukleoseid trutranskriptaseinhibeerder; emtrisitabien is 'n sintetiese nukleosiedanaloog van sitidien en tenofovir DF word in vivo na tenofovir, 'n sikliese nukleosiedfosfonaat (nukleotied) analoog van adenosien 5'- monofosfaat omgeskakel.
Efavirens:
Efavirens is 'n nie-nukleoseid trutranskriptaseinhibeerder (NNTTI) van MIV-1. Dit aktiveteit van efavirens word hoofsaaklik deur reaksiegemiddelde inhibisie van MIV-1 trutranskriptase (TT) gemeidseer. Efavirens inhibeer nie MIV-2 en menslike sellulêre DNA-polimerase α, β, γ en δ nie.
Emtrisitabien:
Emtrisitabien, 'n nukleoseid trutranskriptaseinhibeerder (NTTI), is 'n sintetiese nukleosiedanaloog van sitidien. Selfullere eiseine fosforileer emtrisitabien-5'-trifosfaat. Emtrisitabien-5'-trifosfaat kompeteer met die natuurlike substraat, deoksisitidien-5'-trifosfaat en word in die groeiende virus DNA geïnkorporeer wat tot terminasie van die ketting lei. Emtrisitabien-5'-trifosfaat toon swak inhibisie van soogdier DNA-polimerases α, β, ε en mitochondriale DNA polimerase γ.
Tenofovir disoproksielfumaraat:
Tenofovir DF, 'n NTTI, is 'n sikliese nukleosiedfosfonaat diesteranalog van adenosienmonofosfaat en benodig aanvanklike diesterhidrolise vir omskakeling na tenofovir en daaropvolgende fosforilering deur sellulêre eiseine om tenofovirdifosfaat te vorm. Tenofovirdifosfaat inhibeer die aktiwiteit van MIV-1 TT deur kompetisie met die natuurlike deoksisitidien-5'-trifosfaat te kompeteer. Na inkorporering in DNA, word die DNA-ketting getermineer. Tenofovirdifosfaat toon swak inhibisie teen soogdier polimerases α, β en mitochondriale DNA polimerase γ.

Antivirusaktiwiteit:
Efavirens, emtrisitabien en tenofovir disoproksielfumaraat:
In kombinasiestudies wat die antivirusaktiwiteit in vitro van emtrisitabien en efavirens saam, efavirens en tenofovir saam en emtrisitabien en tenofovir saam gevalueer het, is bykomstige tot sinergistiese antiviruseffekte waargeneem.
Weerstand:
Efavirens, emtrisitabien en tenofovir disoproksielfumaraat:
MIV-isolate met verminderde vatbaarheid vir die emtrisitabien en tenofovir-kombinasie is *in vitro* geselekteer.
Na genotipiese ontleding is die M184V/I en/of K65R aminosuurveranderings in die virus TT geïdentifiseer.
Efavirens:
Kliniese isolate met verminderde vatbaarheid *in vitro* teen efavirens is verkry. Die mees dikwels waargenome aminosuurveranging wat in kliniese studies met efavirens waargeneem is, is K103N (54 %) by monsters met 'n verswakte gevoeligheidsverlies 98, 100, 101, 103, 106, 108, 179, 181, 188, 190, 225, 227 en 230 is waargeneem in pasiente in wie die behandeling met efavirens in kombinasie met ander antiretrovirale middels misluk het. Ander weerstandige mutasies wat dikwels voorkom het L100I (7 %), K101E/Q/R (14 %), V106I (11 %), G190S/T/A (7 %), P225H (18 %) en M230L (11 %).
MIV-isolate met verminderde vatbaarheid vir efavirens as die 290-voudige toename in EC_{50} waarde) het virusig onder seleksie *in vitro* ontwikkel. Genotipiese karakterisering van hierdie viruses het mutasies gedefinieer wat tot enkel aminosuurveranderings L100I of V179D, dubbel veranderings L100I/K103N en dreidubbel veranderings L100I/V179D/Y181C in TT gelei het.
Emtrisitabien:
MIV-isolate wat weerstand teen emtrisitabien bied is *in vitro* geselekteer. Genotipiese ontleding van hierdie isolate het aangetoon dat huitranse by die M184V/I-substitusie in die virus TT ontwikkel het. Geen verminderde vatbaarheid teenoor die MIV TT-geen by kodon 184 geassosieer is. Hierdie mutasie het 'n aminosuurvervanging van meionien deur valien of isoleusien (M184V/I) veroorsaak.
Tenofovir disoproksielfumaraat:
Isolate van MIV-1 wat verminderde vatbaarheid vir tenofovir getoon het is *in vitro* geseleksteer. Hierdie viruse het 'n K65R mutasie in trutranskriptase aangetoon en 'n 2- tot 4-voudige afname in vatbaarheid teen tenofovir getoon.

Kruisweerstand:
Efavirens, emtrisitabien en tenofovir disoproksielfumaraat:
Kruisweerstand is onder NNTTIs uitgewys. Dit is ook bekend dat kruisweerstand onder sekere NTTIs kan ontwikkel. Dat M184V/I met K65R verwerping gaan of nie deur die kombinasie van emtrisitabien en tenofovir geseleksteer is, is ook in somtige MIV-1 stamme uit individue in wie die behandeling met tenofovir of emtrisitabien, en aksaavsuur of didanosien misluk het, waargeneem. Kruisweerstand onder hierdie middels kan daarom in pasiente wat as virus een of beide van hierdie aminosuurveranderings herberg, ontwikkel.

Farmakokinetiese eienskappe:
Efavirens:
Absorpsie:
Piek plasmakonsentrasies van efavirens van 1,6 - 9,1 µM is na 5 ure na 'n enkel dosering van 100 mg tot 1600 mg aan onbesmette vrywilligers bereik. Alhoewel dosisverwante toenames in K_{max} en AOK waargeneem is vir dosisse van tot sowel as 1600 mg, is hierdie toenames kleiner as proporsioneel, wat op 'n afname in absorpsie teen hoer dosisse dui. Plato plasmakonsentrasies word na 8 tot 10 dae bereik.
Verspreiding:
Efavirens is hoogs proteïengebonde (ongeveer 99,5 % tot 99,75 %), hoofsaaklik aan albumien. Konsentrasies in die serebrospinale vloeistof het met 0,26 tot 1,19 % (gemiddeld 0,69 %) van die ooreenstemmende plasmakonsentrasies in MIV-1 besmette individue, aan wie 200 tot 600 mg efavirens een keer per dag vir ten minste een maand toegedien is, gemeisel. Dit is ongeveer die keer meer as die nie-proteïengebonde (vry) fraksie van efavirens in plasma.
Metabolisme:
Efavirens word hoofsaaklik deur die sitochroom P450 sisteem na gehidroksileerde metaboliete gemetaboliseer wat daaropvolgende glukuronidase van hierdie gehidroksileerde metaboliete. Die metaboliete is nie teen MIV-1 aktief nie. CYP3A4 en CYP2B6 is die belangrikste isoënsieme wat in die metabolisme van efavirens betrokke is. Efavirens, deur induksie van P450 ensieme, induseer sy eie metabolisme.
Eliminasie:
Efavirens besit 'n lang terminale halfleeftyd van 52 – 76 ure en 40 - 50 ure na enkel en veelvoudige dosisse, respekteelik. Na die toediening van 'n enkel dosis, is 14 - 34 % van efavirens in die urien (hoofsaaklik as metaboliete) en 16 - 61 % in die feses herwin (hoofsaaklik as onderbmindig).

Emtrisitabien:
Die farmakokinetiese eienskappe van emtrisitabien word in tabel 1 opgesom.
Absorpsie:
Na orale toediening van emtrisitabien (200 mg) word emtrisitabien vinnig geabsorbeer met piek plasmakonsentrasies wat 1 - 2 ure na dosering voorkom.
Verspreiding:
In vitro binding van emtrisitabien aan menslike plasmaproteïene is < 4 % en is afhanklik van die konsentrasie in die reikwydte van 0,02- 200 µg/ml.
Metabolisme en eliminasie:
Na die toediening van radiogemerkte emtrisitabien word ongeveer 86 % in die urien en 13 % as metaboliete herwin. Die metaboliete van emtrisitabien sluit 3'-sulfoksied diastereomere en hul gekoppelde kunjugaat. Dit is bevestig dat kombinasie van twee gemeurdere fitrasie een aktiewe buisekresie geëlimineer. Na 'n enkel dosis emtrisitabien (200 mg) is die plasmahalfleeftyd van emtrisitabien ongeveer 10 ure.

Tenofovir disoproksielfumaraat:
Die farmakokinetiese eienskappe van tenofovir disoproksielfumaraat word in tabel 2 opgesom.
Absorpsie:
Na orale toediening van tenofovir, word maksimum tenofovir serumkonsentrasies na 1,0 ± 0,4 ure bereik.
Verspreiding:
In vitro binding van tenofovir aan menslike plasmaproteïene (< 0,7 %) is onafhanklik van die konsentrasie oor die reikwydte van 0,01 - 25 µg/ml.
Metabolisme en eliminasie:
Ongeveer 70 - 80 % van die intraveneuse dosis van tenofovir word onveranderd in die urien herwin. Tenofovir word deur 'n kombinasie van glomerulêre filtrasie en aktiewe buisekresie geëlimineer. Na 'n enkel dosis tenofovir, is die terminale halfleeftyd van tenofovir ongeveer 17 ure.

Tabel 1: Enkel dosis farmakokinetiese parameters vir emtrisitabien en tenofovir in volwassenes[1]

	Emtrisitabien	Tenofovir
Vastende Orale biobeskikbaarheid[2] (%)	92 (83,1 - 106,4)	25 (NB - 45,0)
Plasma Terminale eliminasiehalfleeftyd[3] (ure)	10 (7,4 - 18,)	17 (12,0 - 25,7)
K_{max}[3] (µg/ml)	1,8 ± 0,72[3]	0,30 ± 0,09
AOK[3] (µg.uur/ml)	10,0 ± 3,12[3]	2,29 ±0,69
CL_R[3] (ml/min)	302 ± 94	1043 ± 115
CL_{tot}[3] (ml/min)	302 ± 89	243 ± 33

[1] NB = Nie beskikbaar
[2] Mediaan (reikwydte)
[3] Gemiddeld ± SA)
[4] Data aangeleid as platowaardes

Effekte van voedsel op orale absorpsie:
TRIEMTA TABLETS is nie in die teenwoordigheid van voedsel gebruiker nie. Die toediening van efavirens tablette met hoë vet maaltye het die gemiddelde AOK en K_{max} van efavirens met 28 % en 79 % respekteelik, verhoog in vergelyking met toediening in die vastende staat. In vergelyking met vastende toediening het die dosering van tenofovir DF en emtrisitabien in kombinasie met hoë vet maaltyd of 'n ligte maaltyd die gemiddelde AOK en K_{max} van tenofovir met 35 % en 15 % respekteelswelk verhoog, sonder om die blootstelling aan emtrisitabien te beïnvloed.

Spesiale populasies:
Pediatriese en bejaarde pasiënte:
Farmakokinetiese studies met tenofovir DF is nie in pediatriese pasiënte (jonger as 18 jaar oud) uitgevoer nie. Effekte van ras en geslag is nie pasiënte gevoer nie. Geen studies met **TRIEMTA TABLETS** is in pediatriese pasiënte vanaf 3 maande tot 17 jaar oud uitgevoer. **TRIEMTA TABLETS** word nie vir pediatriese toediening aanbeveel nie. Die farmakokinetiese van efavirens, emtrisitabien en tenofovir is nie volledig in bejaardes (ouer as 65 jaar) geëvalueer nie (sien **WAARSKUWINGS EN SPESIALE VOORSORGSMAATREËLS**).

Pasiënte met ingekorte nierfunksie:
Efavirens:
Die farmakokinetika van efavirens is nie voldoende in pasiënte met ingekorte lewerfunksie bestudeer nie (sien **WAARSKUWINGS EN SPESIALE VOORSORGSMAATREËLS**).

Emtrisitabien:
Die farmakokinetika van efavirens is nie voldoende in pasiënte met ingekorte lewerfunksie bestudeer nie (sien **WAARSKUWINGS EN SPESIALE VOORSORGSMAATREËLS**).
Farmakokinetiese parameters vir emtrisitabien is verkry in MIV-geïnfekteerde pasiënte met verskillende grade van nierversaking wat nie nodig dialiseer is nie. Die gemiddelde totale liggaamsklaring van emtrisitabien is proporsioneel met die graad van nierversaking, met 'n gemiddelde verlaging in die totale liggaamsklaring van 3 % (gemiddeld 0,03 % van die oorspronklike waarde) in pasiënte met ernstige nierversaking (kreatienienklaring < 30 ml/min). Ongeveer 9 % (reikwydte 6 % tot 15 %) van 'n emtrisitabien dosis is gedurende 'n hemodialisesessie van 3 uur verwyder. Peritoneale dialise het 'n gemiddelde effek van 7 % op die totale liggaamsklaring van emtrisitabien gehad.
Die farmakokinetiese parameters vir emtrisitabien is verkry in MIV-geïnfekteerde pasiënte met verskillende grade van nierversaking wat nie nodig dialiseer is nie. Die gemiddelde totale liggaamsklaring van emtrisitabien is proporsioneel met die graad van nierversaking, met 'n gemiddelde verlaging in die totale liggaamsklaring van 3 % (gemiddeld 0,03 % van die oorspronklike waarde) in pasiënte met ernstige nierversaking (kreatienienklaring < 30 ml/min). Ongeveer 9 % (reikwydte 6 % tot 15 %) van 'n emtrisitabien dosis is gedurende 'n hemodialisesessie van 3 uur verwyder. Peritoneale dialise het 'n gemiddelde effek van 7 % op die totale liggaamsklaring van emtrisitabien gehad.

Tenofovir disoproksielfumaraat:
Die farmakokinetiese parameters vir tenofovir is in MIV-geïnfekteerde pasiënte met verskillende grade van nierversaking wat nie nodig dialiseer is nie, bepaal. Die gemiddelde totale liggaamsklaring van tenofovir is proporsioneel met die graad van nierversaking, met 'n gemiddelde verlaging in die totale liggaamsklaring van 3 % (gemiddeld 0,03 % van die oorspronklike waarde) in pasiënte met ernstige nierversaking (kreatienienklaring < 30 ml/min). Ongeveer 9 % (reikwydte 6 % tot 15 %) van 'n emtrisitabien dosis is gedurende 'n hemodialisesessie van 3 uur verwyder. Peritoneale dialise het 'n gemiddelde effek van 7 % op die totale liggaamsklaring van emtrisitabien gehad.

Etniese groepe:
Farmakokinetiese studies met **TRIEMTA TABLETS** ontwang, waargeneem.
Farmakokinetiese studies met tenofovir DF word wanneer 'n pasiënt bulkyn, naarheid, braking of verwyder is nie bestudeer nie.

Pankreatitis:
Pankreatitis is in sommige pasiënte wat **TRIEMTA TABLETS** ontvang, waargeneem.
Pankreatitis moet oorweg word wanneer 'n pasiënt bulkyn, naarheid, braking of verwydose hoofsaaklik as metaboliete deur die nierfunksie geëlimineer word. Alhoewel dosisverwante toenames in K_{max} en AOK waargeneem is vir dosisse van tot sowel as 1600 mg, is hierdie toenames kleiner as proporsioneel, wat op 'n afname in absorpsie teen hoer dosisse dui. Plato plasmakonsentrasies word na 8 tot 10 dae bereik.

Pediatriese en adolescent gebruik:
Die veiligheid en effektiviteit van **TRIEMTA TABLETS** is nie in kinders en adolescents jonger as 18 jaar vasgestel nie, daarom **TRIEMTA TABLETS** word nie in hierdie ouderdomsgroep aanbeveel nie.

Gebruik in bejaardes:
Kliniese studies van vastedosis kombinasies van emtrisitabien en tenofovir, soos in **TRIEMTA TABLETS**, het nie genoeg individue van ouer as 65 jaar ingesluit om vas te stel of hulle anders as jonger individue reageer nie. Oor die algemeen behoort sorg aan die dag gelê te word wanneer dosisse vir bejaarde pasiënte geseleksteer word, wat groter frekwensie van verminderde lewer-, nier-, en hartfunksie, asook meegaande siekte of medisinale terapie, behoort in gedagte gehou te word.

Effek op die toetsmetode vir bestuurvermoë:
TRIEMTA TABLETS kan duiseligheid, ingekorte konsentrasievermoë en/of tonsiferigheid lei. Pasiente behoort aangeraai te word hulle hierdie simptome ontwikkel, hulle potensieel gevaarlike take soos om te bestuur en met masjinerie te werk, moet vermy.

INTERAKSIES:
Efavirens:
In vivo is daar aangetoon dat efavirens CYP3A4 en CYP2B6 induseer. Ander middels wat substrate van CYP3A4, 2B6 of beide hierdie isoënsieme is, behoort gedoseer te word met inagneming van hierdie bewetenskap. *In vitro* studies het aangetoon dat efavirens 2C9 en 2C19 isoënsieme in die reikwydte van waargeneme plasmakonsentrasies van efavirens geïnhiseer het. Die moontlikheid bestaan dat efavirens saam met medisyne wat hoofsaaklik deur hierdie ensieme gemetaboliseer word, kan tot verminderde klaring van hierdie medisyne en verhoogde blootstelling lei.
Vroeë opsporing en behandeling van die lewerversaking en die ondersteunende sisteng van **TRIEMTA TABLETS** of efavirens word aanbeveel. Pasiente met bestaande lewersiekte, insluitende chronies aktiewe hepatitis toon 'n toename in die frekwensie van lewerfunksie-abnormaliteite gedurende kombinasie antiretrovirale terapie en behoort gemonitor te word soos standaard van sorg. As daar bewyse van verslegtende lewer siekte is, behoort tydelike of permanente staking van behandeling oorweg te word.

Daar is 'n paar bewyse dat efavirens geassosieer word met drie kliniese patologiese patrone van gerwesnstelsel disfunksie: psigotiese simptome, verlenging van normale elektrokardiografie babasc/kinders, behoort klinies en laboratoriumopvolg te ondergaan en behoort in die geval van relevante tekens of simptome ten volle vir misbruikende disfunksie ondersoek te word.

Pankreatitis:
Pankreatitis is in somnige pasiente wat **TRIEMTA TABLETS** ontvang, waargeneem.
Pankreatitis moet oorweg word wanneer 'n pasiënt bulkyn, naarheid, braking of verwydose hoofsaaklik as metaboliete deur die nierfunksie geëlimineer word. Alhoewel dosisverwante toenames in K_{max} en AOK waargeneem is vir dosisse van tot sowel as 1600 mg, is hierdie toenames kleiner as proporsioneel, wat op 'n afname in absorpsie teen hoer dosisse dui. Plato plasmakonsentrasies word na 8 tot 10 dae bereik.

Pediatriese en adolescent gebruik:
Die veiligheid en effektiviteit van **TRIEMTA TABLETS** is nie in kinders en adolescents jonger as 18 jaar vasgestel nie, daarom **TRIEMTA TABLETS** word nie in hierdie ouderdomsgroep aanbeveel nie.

Gebruik in bejaardes:
Kliniese studies van vastedosis kombinasies van emtrisitabien en tenofovir, soos in **TRIEMTA TABLETS**, het nie genoeg individue van ouer as 65 jaar ingesluit om vas te stel of hulle anders as jonger individue reageer nie. Oor die algemeen behoort sorg aan die dag gelê te word wanneer dosisse vir bejaarde pasiënte geseleksteer word, wat groter frekwensie van verminderde lewer-, nier-, en hartfunksie, asook meegaande siekte of medisinale terapie, behoort in gedagte gehou te word.

Effek op die toetsmetode vir bestuurvermoë:
TRIEMTA TABLETS kan duiseligheid, ingekorte konsentrasievermoë en/of tonsiferigheid lei. Pasiente behoort aangeraai te word hulle hierdie simptome ontwikkel, hulle potensieel gevaarlike take soos om te bestuur en met masjinerie te werk, moet vermy.

Tabel 3:
Bevestigde en moontlike interaksies tussen middelinteraksies:
Verandering in dosis of regimen kan op grond van middelinteraksie-studies of voorspelde interaksies gebaseer word.

Meegaande medisyne Kias: Naam van die medisyne	Effek	Kliniese kommentaar
Antiretrovirusmiddels		
Proteaseinhibeerder:	↓ Amprenavir-konsentrasie	Efavirens besit die potensiaal om die serumkonsentrasies van amprenavir te verminder
Amprenavir		
Proteaseinhibeerder: Fosamprenavirkalsium	↓ Amprenavir-konsentrasie	Fosamprenavir (ongesteraag): Toegaslike dosisse van fosamprenavir en **TRIEMTA TABLETS** kan ongun vir veiligheid en effektiwiteit is nie vasgestel nie. Fosamprenavir/ritonavir: 'n Addisionele 100 mg per dag (300 mg totaal) ritonavir word aanbeveel wanneer **TRIEMTA TABLETS** saam met fosamprenavir een keer per dag toegedien word. Wanneer 'n verhoogde dosis ritonavir nodig is nodig wanneer **TRIEMTA TABLETS** saam met fosamprenavir/ritonavir toegedien word nie. Geen verandering in die dosisse van fosamprenavir of ritonavir word aanbeveel nie. Hierdie kombinasie is nie wanneer **TRIEMTA TABLETS** bestuteer nie.
Cholesterol: Die monitering van cholesterol- en triglieriedvlakke behoort in pasiënte wat met **TRIEMTA TABLETS** behandel word, oorweg te word (sien **NEWE-EFFEKTE**).		
Proteaseinhibeerder: Atasanavir † tenofovir konsentrasies	↓ atasanavir konsentrasies ↑ atasanavir konsentrasies	Gelyktydige toediening van atasanavir en tenofovir DF word nie aanbeveel nie. Indien beenbanmateriaal verminderd word, behoort hoogspesifieke konsultasie verky te word.
Beeneffekte: **TRIEMTA TABLETS** verminder die beenmineraaldigtheid. Beenmonitering behoort in MIV-besmette pasiënte met 'n geskiedenis van patologiese beenbreuke of 'n risiko van osteopenie, oorweeg te word. Alhoewel die effek van supplementering van kalsium en vitamien D nie bestudeer is nie, kan die supplementering vir alle pasiënte van voordeel wees. Indien beenabnormaliteite vermoed word, behoort bespoekte konsultasie verky te word.		
Osteonekrose: Alhoewel die etiologie as multifaktoriaal beskou word (insluitende die gebruik van kortikosteroïede, alkoholmisbruik, ernstige immunosuppressie, hoër liggaamsmassa-indeks), is gevalle van osteonekrose aangemeld, veral in pasiënte met gevorderde MIV-siekte en/of langtermyn blootstelling aan kombinasie antiretrovirale terapie (kART). Pasiënte behoort gevra te word oor enige gewrigspyn, -stywheid, -gesiggetheid of moeite met bewegeng ondervind.		
Proteaseinhibeerder: Indinavir	↓ indinavir konsentrasies	Die optimale indinavir dosis, wanneer in kombinasie met efavirens toegedien word, is nie bekend nie. Verhoging van indinavir dosis na 1000 mg elke 8 ure kompenseer nie vir die toename in die metabolisme van indinavir wanneer efavirens.
Proteaseinhibeerder: Lopinavir/ritonavir	↓ lopinavir konsentrasie ↓ ritonavir konsentrasie ↑ tenofovir konsentrasies	In 'n toename in die dosis van lopinavir/ritonavir tot 5 (6 tablette) een keer per dag kan sodanige verhoging in lopinavir konsentrasies word met efavirens in pasiente wat voorheen behandelingsnaïef was, maar 'n afname in die vatbaarheid teenoor lopinavir nie. **Pasiënte behoort vir tenofovir verwante newe-effekte** gemonitor te word. **TRIEMTA TABLETS behoort in pasiënte wat tenofovir** gebruik, as alternatief vir ander kombinasies met efavirens gebruik te word. Indien nodig, kan inisiasie van antiretrovirale terapie gevorderd word. Pasiënte behoort vir tenofovir verwante newe-effekte gemonitor te word. Tenofovir DF gassiseer met lewerafbraking en verhoging van ALAT. Pasiënte behoort vir tenofovir verwante newe-effekte gemonitor te word. Tenofovir DF en didanosien behoort met sorg voorsien te word as gelyktydige toediening van tenofovir DF en didanosien verhoogde sistemiese didanosien blootstelling veroorsaak. Verlaagde didanosien dosisse word aanbeveel. Die moontlikheid bestaan dat gelyktydige toediening van tenofovir DF en didanosien verhoogde sistemiese didanosien blootstelling veroorsaak. Verlaagde didanosien dosisse word aanbeveel. Die moontlikheid bestaan dat gelyktydige toediening van tenofovir DF en didanosien verhoogde sistemiese didanosien blootstelling veroorsaak. Verlaagde didanosien dosisse word aanbeveel.
Immuunherinstelling en inflammatoriese sindroom: Immuunherinstelling en inflammatoriese sindroom (IRIS) is kimmunopatologiese respons wat vlug op die vinnige herstel van immuunfunksie gerapporteer is in pasiente wat met beide bestaande opportunistiese infeksies en asimptomatiese opportunistiese infeksies, sowel as fisiologiese immuunfunksie, behandel is. Hierdie reaksies word gedink om die gevolg te wees van die verbetering van die immuunrespons, wat die liggaam in staat stel om die patogene wat 'n asimptomatiese opportunistiese infeksie